### **Computer Graphics**

### The Human Visual System (HVS)

**Philipp Slusallek** 


# Light

### Electromagnetic (EM) radiation

- From long radio waves to ultra short wavelength gamma rays

#### Visible spectrum: ~400 to 700 nm (all animals)


- Likely due to development of early eyes in water
  - Only very small window that lets EM radiation pass though



# **Plenoptic Function**

#### Physical model for light

- Wave/particle-dualism
  - Electromagnetic radiation wave model
  - Photons:  $E_{ph} = hv \rightarrow$  particle model & ray optics (h: Planck constant)
- Plenoptic function defined at any point in space



### **Radiometric Units**

Specification	Definition	Symbol	Unit	Quantity
Energy		Q <sub>e</sub>	[J = W ⋅s] (joule)	Radiant energy
Power, flux	dQ/dt	Φ <sub>e</sub>	[W = J/s] (watt)	Radiant flux
Flux density	dQ/dAdt	E <sub>e</sub>	[W/m <sup>2</sup> ]	Irradiance
Flux density	dQ/dAdt	B <sub>e</sub>	[W/m <sup>2</sup> ]	Radiosity
Intensity	dQ/d∞dt	l <sub>e</sub>	[W/sr]	Radiant intensity
	dQ/dAd∞dt	L <sub>e</sub>	[W/(m² ·sr)]	Radiance


# Photometry

#### Equivalent units to radiometry

- Weighted with luminous efficiency function V( $\lambda$ )
- Considers the spectral sensitivity of the human eye
  - Measured across different humans
- Spectral or (typically) "total" units
  - Integrate over the entire spectrum and deliver a single scalar value

$$\Phi_{v} = K_{m} \int V(\lambda) \Phi_{e}(\lambda) d\lambda$$
$$K_{m} = 680 \, lm/W$$

- Simple distinction (in English!):
  - Names of radiometric quantities contain "radi"
  - · Names of photometric quantities contain "lumi"



function

### **Photometric Units**

Specification	Definition	Symbol	Unit	Quantity
Energy		Q <sub>v</sub>	[T = Im ⋅s] (talbot)	Luminous energy
Power, flux	dQ/dt	Φν	[lm = T/s] (lumen)	Luminous flux (e.g. emitted power of lamp)
Flux density	dQ/dAdt	Ev	$[lx = lm/m^2]$ (lux)	Illuminance (e.g. illumination on desk)
Flux density	dQ/dAdt	B <sub>v</sub>	$[lx = lm/m^2]$ (lux)	Luminosity (e.g. reflection off desk)
Intensity	dQ/dωdt	I <sub>v</sub>	[cd = lm/sr] (candela)	Luminous intensity (e.g. intensity of a point light)
	dQ/dAd∞dt	L <sub>v</sub>	[lm/(m² ⋅sr)] (nits)	Luminance (e.g. brightness of a monitor)


With luminous efficiency function weighted units

# **Illumination: Examples**


#### Typical illumination intensities

Light source	Illuminance [lux]	
Direct solar radiation	25,000 - 110,000	
Day light	2,000-27,000	
Sunset	1 – 108	
Moon light	0.01 – 0.1	
Starry night	0.0001 - 0.001	
TV studio	5,000 - 10,000	
Shop lighting	1,000 - 5,500	
Office lighting	200 – 550	
Home lighting	50 – 220	
Street lighting	0.1 – 20	


# Luminance Range



# Contrast (Dynamic Range)




# High Dynamic Range (HDR)



- How to display computed/measured HDR values on an LDR device ?
  - Tone mapping ( $\rightarrow$  RIS course)


# Percept. Effects: Vision Modes



#### • Simulation requires:

- Control over color reproduction
- Local reduction of detail visibility (computationally expensive)


### Visual Acuity and Color Perception



Simulation, (c) Cornell

# Percept. Effects: Temp. Adaptati.


Adaptation to dark much slower



I sudden change in illumination


- Simulation requires:
  - Time-dependent filtering of light adaptation

### **HVS - Relationships**



### Human Visual System


- Physical structure well established
- Percept. behavior complex & less understood process




Optic chiasm

# **Optical Chiasm**


- Right half of the brain operates on left half of the field of view
  - From both eyes!!
- And vice versa
  - Damage to one half of the brain can results in loss of one half of the field of view



### **Perception and Eye**



### **Human Visual Perception**




early vision (eyes)

- Determines how real-world scenes appear to us
- Understanding of visual perception is necessary to reproduce appearance, e.g. in tone mapping


# **Distribution of Rods and Cones**

- High-res. foveal region with highest cone density
- Poisson-disc-like distribution




### Retina

- Receptors on opposite side of incoming light
- Early cellular processing between receptors & nerves
  - Mainly for rods




### Eye as a Sensor

#### Relative sensitivity of cones



### Luminuous Sensitivity Function

• Different for cones (black, diff. studies) & rods (green)



# Eye

#### Fovea (centralis):

- Ø 1-2 visual degrees
- 50,000 cones each of ~ 0.5 arcminutes angle (~2.5 µm wide)
- No rods in central fovea, but three different cone types:
  - L(ong, 64%), M(edium, 32%), S(hort wavelength, 4%)
  - $\Rightarrow$  Varying resolution: 10 arcminutes for S vs. 0.5 arcminutes for L & M
- Mostly linked directly with optical nerves and visual cortex (1:1),
  - 1% of retina area but covers 50% of visual cortex in brain
- Adaptation of light intensity only through cones

#### Periphery:

- 75-150 M. rods: night vision (B/W)
- 5-7 M. cones (color)
- Rods: Response to stimuli by even a single photon (@ 500 nm)
  - 100x better than cones, integrating over 100 ms
- Signals from many rods are combined before linking with nerves
  - Bad resolution, good flickering sensitivity

This is a text in red

This is a text in green

This is a text in blue

This is a text in red

This is a text in green

This is a text in blue

This is a text in red

This is a text in green

This is a test in blue

#### This is a text in green

#### This is a text in blue

This is a text in green

This is a text in blue

This is a text in green

This is a test in blue

#### This is a text in red


This is a text in green

This is a text in red

This is a text in green

This is a text in red

This is a text in green



# **Resolution of the Eye**

#### Resolution-experiments

- Line pairs: eye ~ 50-60 p./degree  $\rightarrow$  resolution of 0.5 arcminutes
- Line offset: 5 arcseconds (hyperacuity)



- Eye micro-tremor: 60-100 Hz, 5 μm (2-3 photoreceptor spacing)
  - Allows to create super-resolution (w/ Poisson pattern)
- Together corresponds to 19" display at 60 cm away from viewer: 3,000<sup>2</sup> without hyperacuity – 18,000<sup>2</sup> pixels with hyperacuity

#### Fixation of eye onto (moving) region of interest

- Automatic gaze tracking, autom. compensation of head movement
- Apparent overall high resolution of fovea


#### Visual acuity increased by

- Brighter objects and high contrast

### **Contrast Sensitivity**

#### Human visual system

- Perception very sensitive to regular structures
- Insensitive against (high-frequency) noise
- Campbell-Robson sinusoidal contrast sensitivity chart




# Luminance Contrast Sensitivity

- Sensitivity: inverse of perceptible contrast threshold
- Maximum acuity at 5 cycles/degree (0.2 %)
  - Decrease toward low frequencies: lateral inhibition
  - Decrease toward high frequencies: sampling rate (Poisson disk)
  - Upper limit: 60 cycles/degree

### Medical diagnosis


- Glaucoma (affects peripheral vision: low frequencies)
- Multiple sclerosis (affects optical nerve: notches in contrast sensitivity)
   Human Contrast Sensitivity




# **Color Contrast Sensitivity**

#### Color vs. luminance vision system

- Similar but slightly different curves
- Higher sensitivity at lower frequencies
- High frequencies less visible
- Image compression
  - Exploit color sensitivity in lossy compr.







# **Threshold Sensitivity Function**


#### • Weber-Fechner law (Threshold Versus Intensity, TVI)

- Perceived brightness varies linearly with log(radiant intensity)
  - E = K + c log l
- Perceivable intensity difference
  - 10 cd vs. 12 cd: ΔL = 2 cd
    20 cd vs. 24 cd: ΔL = 4 cd
    - 30 cd vs. 36 cd:  $\Delta L = 6$  cd





### Weber-Fechner Examples




### Mach Bands

#### "Overshooting" along edges

- Extra-bright rims on bright sides
- Extra-dark rims on dark sides

#### Due to "lateral inhibition"




### Mach Bands

#### "Overshooting" along edges

- Extra-bright rims on bright sides
- Extra-dark rims on dark sides

#### Due to "lateral inhibition"




# Lateral Inhibition

#### Pre-processing step within retina

- Surrounding brightness level weighted negatively
  - A: high stimulus, maximal bright inhibition
  - B: high stimulus, reduced inhibition  $\rightarrow$  stronger response
  - D: low stimulus, maximal dark inhibition
  - C: low stimulus, increased inhibition  $\rightarrow$  weaker response

### High-pass filter

- Enhances contrast along edges
- Differential operator (Laplacian/difference of Gaussian)



# Lateral Inhibition: Hermann Grid

#### Apparent dark spots at perip. crossings


- Weakly if within foveal  $\Omega$  (B): smaller filter extent
- Strongly within periphery (A): larger filter extent


#### Explanation


- Crossings (C): more surround stimulation
  - More inhibition  $\Rightarrow$  weaker response
- Streets (D): less surround stimulation
  - Less inhibition  $\Rightarrow$  greater response


#### Simulation

- Convolution with differential kernel
- Darker at crossings, brighter in streets












### **Some Further Weirdness**



# **High-Level Contrast Processing**




# **High-Level Contrast Processing**



### **Cornsweet Illusion**


Apparent contrast between inner and outer shades



Β


### **Cornsweet Illusion**

- Apparent contrast between inner and outer shades
  - Due to gradual darkening/brightening towards a contrasting edge
  - Causes B to be perceived similarly to A



# **Optical Effects – Veiling Glare**


- Internal scattering/blur of sources of high luminance
- Blur around the bright object makes it appear brighter!





# **Shape Perception**

#### Depends on surrounding primitives

- Size emphasis
- Directional emphasis









### **Geometric Cues**


#### Automatic geometrical interpretation

- 3D perspective
- Implicit scene depth





### Visual "Proofs"




# **HVS: High-Level Scene Analysis**

#### Experience & expectation

- Pictures usually horizontal
- Local cue consistency
  - Eyes and mouth look right, but actually are upside-down







# **HVS: High-Level Scene Analysis**

#### Experience & expectation

- Pictures usually horizontal
- Local cue consistency
  - Eyes and mouth look right, but actually are upside-down








### **Impossible Scenes**

#### • Escher et al.


- Confuse HVS by presenting contradicting visual cues
- Locally consistent but not globally






### Single Image Random Dot Stereograms

- Vergence: Cross eyers to look at the same 3D spot
- Accommodation: Focusing at a particular depth plane



# **SIRDS** Construction


- Assign arbitrary color to pixel p<sub>0</sub> in image plane
- Trace from eye points through p<sub>0</sub> to object surface
- Trace back from object to corresponding other eye
- Assign color at  $p_0$  to intersection points  $p_{1L}$ ,  $p_{1R}$  with image plane
- Trace from eye points through p<sub>1L</sub>,p<sub>1R</sub> to object surface
- Trace back to eyes
- Assign p<sub>0</sub> color to p<sub>2L</sub>, p<sub>2R</sub>
- Repeat until image plane is covered




# **Motion Illusion**

#### Appearance of movement in static image


- Due to cognitive effects of interacting color contrast & shape pos.
- Saccades  $\rightarrow$  diff. in neural signals between dark and bright areas



### **Motion Illusion**



### **Motion Illusion**



# **Negative Afterimages**

- Cones excited by color eventually lose sensitivity
  - Photoreceptors adapt to overstimulation and send a weak signal



# **Negative Afterimages**

- When switching to grey background
  - Colors corresponding to adapted cones remain muted
  - Other freshly excited cones send out a strong signal
  - Same perceived signal as when looking at the inverse color



### **Another Optical Illusion**

• If staring for ~ 15 sec., you may see a giraffe appear

